



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2013

Comparison of age of peak swimming speed in elite backstroke swimmers at national and international level

Kollarz, C ; Knechtle, B ; Rüst, C A ; Rosemann, T

Abstract: Background: Previous studies suggested that the age of peak performance in freestyle swimming was 17 years for women and 19 years for men. Data on the age of peak swimming performance in other swim strokes such as backstroke are lacking. The present study investigated the changes in (i) swimming speed and (ii) age of peak swimming speed across years for elite female and male backstroke swimmers competing at both national and international level. Methods: The changes in age and swimming speed in 50m, 100m and 200m were analyzed for elite backstroke and freestyle swimmers at national level (i.e. Switzerland) and international level (i.e. finalists of World Championships) on 50m long course from 1994 to 2011. Results: For all distances and disciplines, women reached the age of peak swimming speed at a younger age (18-23 years) than men (21-26 years). The age of peak swimming speed in backstroke and freestyle swimmers at national level was 1-2 years younger than in swimmers at international level. The age of peak swimming speed increased significantly ($p < 0.01$) between 1994 and 2011 in 50m backstroke for women from 16 to 22 years and in 50m freestyle for men from 22 to 23 years in swimmers at national level. No changes in the age of peak swimming speed were observed in swimmers at international level for both swim styles across time for both sexes ($p > 0.05$). Swimming speed increased across years in all distances and disciplines except for 50m backstroke in women at international level ($p > 0.05$). Conclusion: Women reached in all disciplines the age of peak swimming speed at a younger age than men. Swimmers at national level were younger than swimmers at international level. Swimming speed increased in all distances and disciplines across years in swimmers competing at both national and international level.

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-92144>

Journal Article

Published Version

Originally published at:

Kollarz, C; Knechtle, B; Rüst, C A; Rosemann, T (2013). Comparison of age of peak swimming speed in elite backstroke swimmers at national and international level. *OA Sports Medicine*, 1(2):19.

Comparison of age of peak swimming speed in elite backstroke swimmers at national and international level

C Kollarz¹, B Knechtle^{1,2*}, CA Rüst¹, T Rosemann¹, R Lepers³

Abstract

Introduction

Previous studies suggested that the age of peak performance in freestyle swimming was ~17 years for women and ~19 years for men. Data on the age of peak swimming performance in other swim strokes such as backstroke are lacking. The present study investigated the changes in swimming speed and age of peak swimming speed across years for elite female and male backstroke swimmers competing at both national and international levels.

Materials and methods

The changes in age and swimming speed in 50 m, 100 m and 200 m were analysed for elite backstroke and freestyle swimmers at national level (i.e. Switzerland) and international level (i.e. finalists of World Championships) on the 50-m long course from 1994 to 2011.

Results

For all distances and disciplines, women reached the age of peak swimming speed at a younger age (~18–23 years) than men (~21–26 years). The age of peak swimming speed in backstroke and freestyle swimmers at national level was ~1–2 years younger than in swimmers at international level. The age of peak swimming speed increased significantly ($P < 0.01$) between 1994 and 2011 in 50-m backstroke for women from

16 to 22 years and in 50-m freestyle for men from 22 to 23 years in swimmers at national level. No changes in the age of peak swimming speed were observed in swimmers at international level for both swim styles across time for both sexes ($P > 0.05$). Swimming speed increased across years in all distances and disciplines except for 50-m backstroke in women at international level ($P > 0.05$).

Conclusion

Women reached in all disciplines the age of peak swimming speed at a younger age than men. Swimmers at national level were younger than swimmers at international level. Swimming speed increased in all distances and disciplines across years in swimmers competing at both national and international levels.

Introduction

Several studies analysed the change of endurance performance with increasing age^{1–3}. For endurance sports such as running, swimming, triathlon or rowing, peak performance was described at the age of ~35 years with a linear decline in performance until the age of ~70 years^{2,4,5}. After the age of ~70 years, the performance declined exponentially^{1–3,5}.

For swimmers, the age-related performance decline in freestyle swimming showed differences regarding the distance and the sex, with a greater decline in long-distance performances such as 1,500-m freestyle and for women compared to men^{2,3}. Schulz and Curnow⁶ reported that women generally achieved peak swimming performance at a younger age compared to men. Literature, however, is discordant regarding the age of peak freestyle swimming

performance in relation to different race distances. For freestyle swimmers, the age of peak swimming performance was described at ~21 years⁷. Fairbrother⁸ reported that male freestyle swimmers achieved the age of peak performance for 50 m in their late 20s and early 30s. For 1,500-m freestyle, the age of peak swimming performance was achieved at ~18 years in contrast to the 50-m distance where the age of peak swimming speed was documented at ~23 years⁷. Schulz and Curnow⁶ described a different development with a decline in the age of peak swimming performance with increasing race distance. Tanaka and Seals³ claimed that men reached their fastest times for 1,500-m freestyle swimming between 25 and 40 years of age, whereas women achieved the fastest 1,500-m freestyle swimming times at the age of ~30–35 years. In contrast, peak performance in 50-m freestyle swimming was attained at the age of 20–30 years in both men and women³.

In contrast to these disparate findings in swimming, data analyses of running showed an increase in the age of peak performance with increasing race distance^{6,9,10}. The age of peak running performance has been investigated in 100-m and 200-m short-distance running⁶, 800-m and 1,500-m middle-distance running⁶, half-marathon running⁹, marathon running^{6,9} and ultra-marathon running¹⁰.

Analysing the development of swimming performances of women compared to men, women's performances constantly improved^{6,11}. During the last century, a decline in the sex difference in performance for all

*Corresponding author
Email: beat.knechtle@hispeed.ch

¹ Institute of General Practice and for Health Services Research, University of Zurich, Zurich, Switzerland

² Gesundheitszentrum St. Gallen, St. Gallen, Switzerland

³ INSERM U1093, Faculty of Sport Sciences, University of Burgundy, Dijon, France




Figure 1: Change in age (panels A, C, E) and swimming speed (panels B, D, F) of the national backstroke swimmers for 50 m (panels A and B), 100 m (panels C and D) and 200 m (panels E and F) with sex difference across the years.

swim disciplines from 22.9% in 1916 to 8.9% in 1979 was documented¹¹. Buhl et al.¹² analysed the change in swimming speed and sex differences in swimming speed for the 200-m and 400-m freestyle and medley from 1994 to 2011. Athletes of both sexes showed significant improvements in all disciplines across years with an increase of sex difference except for the 200-m medley. Except for Buhl

et al.¹², recent data about sex difference in swimming performance are lacking, and it would be interesting to investigate the development of sex differences in swimming performance in the last 30 years. Furthermore, Schulz and Curnow⁶ investigated the age of peak swimming performance for freestyle, but data about different strokes such as backstroke were not considered.

Therefore, the aim of this study was to examine the changes in swimming speed and age of peak swimming speed for both elite male and female swimmers at national and international levels in backstroke and freestyle from 1994 to 2011. There is only a small amount of literature about backstroke swimming in contrast to freestyle swimming; moreover, literature about freestyle swimming includes a great variety of different freestyle swim events. Therefore, there are limitations in the comparability of literature of swimming events. Hence, we decided to compare backstroke with freestyle as internal standard. We hypothesised that (i) the age of peak swimming speed of both elite Swiss swimmers and finalists at World Championships would decrease and swimming speed would increase over time for both women and men and (ii) the age of peak swimming speed would be lower for women compared to men for both backstroke and freestyle. To test this hypothesis, we analysed swimming speed of the top swimmers at national level (i.e. annual Swiss high score list)¹³ and international level (i.e. finalists at World Championships which occurs every 4 years)^{14,15} and compared the age the top 10 swimmers across years in backstroke and freestyle swimming for 50-m, 100-m and 200-m, respectively.

Materials and methods

All procedures used in the study met the ethical standards of the Swiss Academy of Medical Sciences¹⁶ and were approved by the Institutional Review Board of Kanton St. Gallen, Switzerland, with a waiver of the requirement for informed consent of the participants, given the fact that the study involved the analysis of publicly available data.

Data sampling and data analysis

The data set from this study was obtained for national swimmers from

Table 1 Age of peak backstroke swimming speed of national top 10 swimmers from 1994 to 2011 for men and women

	Age 1994	Age 2011	Signifi- cance	Sex differ- ence 1994	Sex differ- ence 2011	Signifi- cance
50-m backstroke						
Women	16 ± 2	22 ± 4	<i>P</i> < 0.01	28.19	2.31	<i>P</i> = 0.01
Men	23 ± 5	22 ± 3	<i>P</i> > 0.05			
100-m backstroke						
Women	18 ± 4	20 ± 3	<i>P</i> > 0.05	12.56	11.11	<i>P</i> > 0.05
Men	21 ± 4	22 ± 3	<i>P</i> > 0.05			
200-m backstroke						
Women	18 ± 3	18 ± 3	<i>P</i> > 0.05	11.59	14.92	<i>P</i> > 0.05
Men	21 ± 2	21 ± 3	<i>P</i> > 0.05			
Age is presented in years as mean (±SD); sex difference is presented in % as mean (±SD).						

Age is presented in years as mean (±SD); sex difference is presented in % as mean (±SD).

the website of the Swiss Swimming Federation¹³ and for international swimmers from the International FINA-Ranking^{14,15}. All swimmers from the Swiss swimming high score list between 1994 and 2011 and the finalists of the World Championships of the International FINA-Ranking on the 50-m long course were analysed regarding swimming speed, sex and age.

Data were available from 1994 to 2011 for 207,852 athletes at national level and 369 athletes at

international level. From the athletes competing at national level, the annual fastest race time and the race times of the annual top 10 athletes for the distances of 50-m, 100-m and 200-m in backstroke and freestyle were determined for both women and men. For the swimmers competing at international level, the eight finalists were included. When there were less than 10 male or female participants for a specific discipline in a year at national level, only the best annual performance was analysed.

Table 2 Age of peak backstroke swimming speed of international FINA World Championship swimmers from 1994 to 2011 for men and women

	Age1994	Age2011	Significance	Sex difference 1994	Sex difference 2011	Significance
50-m backstroke						
Women*	22 ± 4	23 ± 4	<i>P</i> > 0.05	16.44	13.50	<i>P</i> > 0.05
Men*	23±2	26 ± 1	<i>P</i> > 0.05			
100-m backstroke						
Women	21 ± 2	22 ± 4	<i>P</i> > 0.05	13.50	18.63	<i>P</i> > 0.05
Men	24 ± 1	24 ± 3	<i>P</i> > 0.05			
200-m backstroke						
Women	21 ± 3	19 ± 2	<i>P</i> > 0.05	15.01	15.37	<i>P</i> > 0.05
Men	23 ± 2	23 ± 4	<i>P</i> > 0.05			

Age is presented in years as mean (±SD); sex difference is presented in % as mean (±SD). *Data available from 2001.

In order to compare the swimming performance between the disciplines and distances, race times were transformed to swimming speed (m/s) using the equation [race distance in metres]/[race time in seconds]. To compare male and female performances, the sex difference was calculated using the equation ([swimming speed in women] – [swimming speed in men])/[swimming speed in men] × 100, where sex difference was calculated for every pairing of equally placed athletes (e.g. between male and female first place, between male and female second place, etc.) before calculating the mean value and standard deviation (SD) of all the pairings. In order to facilitate reading, all gender differences were transformed to absolute values before analysing. No athlete was included twice or several times in the same year because Swiss Swimming Federation lists only the best performance per year of an athlete in the same year.

Statistical analysis

In order to increase the reliability of data analyses, each set of data was tested for normal distribution as well as for homogeneity of variances in advance of statistical analyses. Normal distribution was tested using a D'Agostino and Pearson omnibus normality test and homogeneity of variances was tested using a Levene's test. To find significant changes in a variable across years, single linear regression analysis was used. Two-way analysis of variance (ANOVA) with Bonferroni post-hoc analysis was used to analyse differences between the annual top 10 women and men. Statistical analyses were performed using IBM SPSS Statistics (Version 19 and 21, IBM SPSS, IL, USA) and GraphPad Prism (Version 5 and 6, GraphPad Software, CA, USA). Significance was accepted at $P < 0.05$ (two sided for t -tests). Data in the text are given as mean ± SD.

Licensee OA Publishing London 2013. Creative Commons Attribution License (CC-BY)

FOR CITATION PURPOSES: Kollarz C, Knechtle B, Rüst CA, Rosemann T, Lepers R. Comparison of age of peak swimming speed in elite backstroke swimmers at national and international level. OA Sports Medicine 2013 Oct 01;1(2):19.




Figure 2: Change in age (panels A, C, E) and swimming speed (panels B, D, F) of the international backstroke swimmers for 50 m (panels A and B), 100 m (panels C and D) and 200 m in backstroke (panels E and F) with sex difference across the years.

Results

Change in the age of peak swimming speed across years

In backstroke, women at national level achieved the age of peak swimming speed at a younger age (16–22 years) (Table 1, Figure 1A, 1C, 1E) compared to women at international level (19–23 years) (Table 2, Figure 2A, 2C, 2E). Only in 50-m backstroke in athletes competing at national level, a change in the age of peak

swimming speed was observed from 16 to 22 years (Table 1, Figure 1A).

The age of peak swimming speed in backstroke was older for men than for women. Furthermore, the age of peak swimming speed was older for swimmers at international level (23–26 years) (Table 2, Figure 2A, 2C, 2E) than in swimmers at national level (21–23 years) (Table 1, Figure 1A, 1C, 1E). However, in backstroke, no changes in the age of peak swimming

speed were observed across years. The sex difference in swimming speed remained stable across the years, with the exception of 50-m backstroke of national swimmers with a decline from 28.2% to 2.3% across years (Table 1, Figure 1A).

In freestyle, no changes in the age of the fastest swimming speed were observed for both sexes in all distances neither on national (Table 3, Figure 3A, 3C, 3E) nor on international (Table 4, Figure 4A, 4C, 4E) level. Only in the 50-m freestyle of men competing at national level (Table 3, Figure 3A), a significant decline from 23 to 22 years was observed. However, again the age of the highest swimming speed was lower for swimmers competing at national level (women 18–21 years; men 21–23 years) compared to athletes competing at international level (women 19–25; men 22–25 years). While the sex differences in swimming speed declined significantly in all distances in athletes at national level (mean from 16.3% to 8.8%) (Table 3, Figure 3A, 3C, 3E), the sex difference in swimming speed of athletes competing at international level remained unchanged across years (mean 17.1% to 14.3%; $P > 0.05$) (Table 4, Figure 4A, 4C, 4E).

Change in swimming speed across years

The swimming speed increased across years, with the exception of the 50-m backstroke for women competing at international level (Table 5). Women showed a smaller improvement in swimming speed than men, with a greater improvement in backstroke than in freestyle. However, swimmers competing at national level (Tables 6 and 7) showed greater changes in swimming speed compared to swimmers competing at international level (Tables 5 and 8). The change in swimming speed depending on distance for women showed an improvement with a range from 4.4% to 8.5% in backstroke and from 2.4% to 2.8% in freestyle in swimmers

Table 3 Age of peak freestyle swimming speed of national top 10 swimmers from 1994 to 2011 for men and women

	Age 1994	Age 2011	Significance	Sex difference 1994	Sex difference 2011	Significance
50-m freestyle						
Women	19 ± 2	20 ± 3	P > 0.05	18.53	12	P < 0.01
Men	23 ± 4	22 ± 3	P < 0.01			
100-m freestyle						
Women	18 ± 2	21 ± 4	P > 0.05	18.92	9.09	P = 0.04
Men	22 ± 4	23 ± 3	P > 0.05			
200-m freestyle						
Women	19 ± 3	20 ± 2	P > 0.05	11.48	5.39	P < 0.01
Men	21 ± 2	22 ± 4	P > 0.05			
Age is presented in years as mean (±SD); sex difference is presented in % as mean (±SD).						

Age is presented in years as mean (±SD); sex difference is presented in % as mean (±SD).

competing at national level. Women competing at international level showed a range from 3.3% to 3.7% in backstroke and from 1.8% to 3.0% in freestyle. Men competing at national level, however, showed a range of improvement in swimming speed from 4.7% to 9.4% in backstroke and from 4.4% to 5.3% in freestyle, while male athletes competing at international level showed a range from 3.0% to 5.6% in backstroke and from 2.7% to 3.5% in freestyle. However, due to a greater improvement in swimming speed in men compared to women, the sex difference in swimming speed increased in the 100-m backstroke (Table 6, Figure 1D) and all freestyle distances of swimmers at national level (Table 7, Figures 2 B, 2D, 2F). In swimmers at international level, the sex differences in swimming speed remained unchanged across years (Tables 5 and 8, Figure 3B, 3D and Figures 4B, 4D, 4F).

Discussion

The aim of this study was to examine the changes in swimming speed and the age of peak swimming speed for both elite male and female swimmers at national and international levels in backstroke versus freestyle from 1994 to 2011. According to our

hypothesis, women achieved peak swimming speed at a younger age than men in both disciplines and all distances. During the studied period, peak swimming speed improved at a greater extent for men than for women in all distances and both strokes on both national and international levels.

Women achieved the age of peak swimming speed at a younger age than men

A first important finding was that women achieved peak swimming speed in freestyle and backstroke at a

younger age than men. In all distances, women reached peak swimming speed at ~18 years in backstroke and at ~19 years in freestyle. For men, peak swimming speed in 50 m was achieved at ~20 years in backstroke and at ~22 years in freestyle and at 21 years in 100 m for both disciplines.

These findings are in line with Schulz and Curnow⁶ investigating the age of peak performances in different sport events including different freestyle swimming events. However, in the present study, athletes of both sexes achieved peak swimming speed in freestyle and backstroke at higher ages compared to previous data. Schulz and Curnow⁶ found the age of peak swimming speed in freestyle for women at ~17 years and for men at ~19 years, respectively. A potential explanation for these different findings might be the analysed period of time. Schulz and Curnow⁶ analysed data from 1896 to 1980, while our results are based on a data from 1994 to 2011. Schulz and Curnow⁶ separated their data into a period before and after World War II. Men showed a decline in the age of peak swimming performance after World War II while women showed an increase after World War II. These

Table 4 Age of peak freestyle swimming speed of international FINA World Championship swimmers from 1994 to 2011 for men and women

	Age 1994	Age 2011	Signifi- cance	Sex differ- ence 1994	Sex differ- ence 2011	Signifi- cance
50-m freestyle						
Women	20 ± 4	25 ± 5	P > 0.05	16.03	14.44	P > 0.05
Men	23 ± 2	25 ± 3	P > 0.05			
100-m freestyle						
Women	20 ± 4	24 ± 3	P > 0.05	15.48%	13.08	P > 0.05
Men	23 ± 3	24 ± 3	P > 0.05			
200-m freestyle						
Women	19 ± 3	21 ± 2	P > 0.05	19.79%	15.42	P > 0.05
Men	22 ± 3	24 ± 3	P > 0.05			
Age is presented in years as mean (±SD); sex difference is presented in % as mean (±SD).						

Age is presented in years as mean (±SD); sex difference is presented in % as mean (±SD).

Licensee OA Publishing London 2013. Creative Commons Attribution License (CC-BY)

FOR CITATION PURPOSES: Kollarz C, Knechtle B, Rüst CA, Rosemann T, Lepers R. Comparison of age of peak swimming speed in elite backstroke swimmers at national and international level. OA Sports Medicine 2013 Oct 01;1(2):19.




Figure 3: Change in age (panels A, C, E) and swimming speed (panels B, D, F) of the national freestyle swimmers for 50 m (panels A and B), 100 m (panels C and D) and 200 m (panels E and F) with sex difference across the years.

changes across the last century can be explained by geopolitical differences and historical developments¹⁷ on the one hand and by developments in training techniques and improvement in swimming techniques related to a greater knowledge of biological and physiological processes¹⁸ on the other hand.

Swimming performances are related to specific body profile markers¹⁹. Therefore, changes in the age of fastest swimming speed can be

related to changes in anthropometric pattern across time. A variety of studies showed an increment on both body height and body weight among children over the last century in different countries such as Japan^{20,21}, Turkey²², Brazil²³, Poland²⁴ and China^{25,26}. This phenomenon seemed to be related to environmental and socioeconomic factors such as changes in optimised nutrition and improvement in health and hygienic systems^{20,26}.

Furthermore, specific body profile markers of swimmers such as body height, length of extremities and body weight were attributed to predict swimming performances in different swim events already at preadolescent age¹⁹. While sprint and backstroke swimmers were taller, heavier and showed longer extremities, short and light swimmers showed a better performance in long-distance and butterfly events¹⁹. Charles and Bejan²⁷ analysed male champions in 100-m freestyle swimming and 100-m dash running from 1912 to 2008. They described that during the last century, male champion swimmers became taller and heavier and achieved higher swimming speeds by producing more power due to optimal locomotion conditions²⁷.

Due to differences in body compositions of men and women, swimming performances of the sexes are related to different anthropometric markers. Geladas et al.²⁸ analysed predictor variables in freestyle swimming performance for men and women. For male swimmers, upper-extremity length, horizontal jump and grip strength were significant predictors for performances in freestyle swimming, whereas body height, upper extremity and hand length, shoulder flexibility and horizontal jump were value predictors for female swimmers²⁸. Based on these facts, sex differences in swimming performance seemed to be influenced by the earlier onset of maturation in women compared to men^{29,30}. These differences in development effects on the one hand bone growth and on the other hand the development and distribution of body fat.

Regarding bone growth, girls reach the maximum of all bone variables at younger age than boys³¹. Martin et al.²⁹ described a difference of approximately 2 years between the growth of metacarpal bones between girls and boys. Compared to an adult bone mineral content and volumetric den-

Table 5 Changes in swimming speed in backstroke across from 1994 to 2011 for international FINA World Championship swimmers

	Changes (m/s)	Changes (%)	Significance	Sex difference 1994	Sex difference 2011	Significance
50-m backstroke						
Women	+0.04	+2.3	P > 0.05	10.28	11.50	P > 0.05
Men	+0.08	+4	P = 0.03			
100-m backstroke						
Women	+0.06	+3.7	P < 0.01	9.58	10.07	P > 0.05
Men	+0.10	+5.6	P<0.01			
200-m backstroke						
Women	+0.05	+3.3	P = 0.01	9.46	9.14	P > 0.05
Men	+0.05	+3	P < 0.01			
Changes in swimming speed are presented in m/s and %.						

Changes in swimming speed are presented in m/s and %.

sity, boys reach approximately 86% while girls show 93% at an age of 17 years³¹. However, the performance of swimmers between 14 and 16 years in 100-m freestyle was partly predictable by anthropometry and physical capacity with lower value for boys and girls²⁸.

Besides different changes of the skeleton during puberty, boys and girls show a different development in body fat and muscle mass. The change in both the proportion and the distribution of body fat seemed to influence swimming performance^{32–34}. During puberty, girls develop more fat mass compared to boys, while both gain more muscle mass³⁵. At the age of ~11 years, boy sprint swimmers presented 9.4% fat and girl sprint swimmers 12.7% fat³⁴. In the last stages of and after puberty, women obtain more fat mass³⁰. Adult female swimmers have a per cent body fat of 14–19% than male swimmers of 5–10%³⁶. The locations of body fat and body mass affect the centres of buoyancy, and this influences kick performance and swimming performances respectively³⁷. Especially, women have proportionally more fatty tissue located caudally compared to men, and therefore centre of buoyancy is different for the

sexes. Special training programs may help to reduce the proportion of body fat especially for women to improve their swim performances³⁴.

Sex difference in swimming performance

Men achieved a higher swimming speed in all distances and disciplines than women, with higher swimming speeds in freestyle compared to backstroke. Furthermore, swimming speed declined with increasing distance for all distances and disciplines on national and international levels.

For both sexes and all disciplines and distances, the swimming speeds of the international athletes were higher than of the national swimmers. This finding is in line with existing literature^{12,38}.

Swimming speed depends on a variety of different patterns. Besides anthropometric factors as mentioned, different physiological parameters and swimming techniques have an impact on swim speed³⁹. Physiological pattern such as metabolic responses and the ability of the body to adapt have an effect on swimming speed. The energy cost of a performance is related to the oxygen consumption, cardiopulmonary response and metabolic adaptations⁴⁰. Analysing the energy cost of different swimming styles, freestyle has a lower value than backstroke⁴¹. Men have a higher $\dot{V}O_{2\max}$ compared to women, but this difference disappears when normalising to leg mass⁴⁰. Fomin et al.⁴⁰ analysed the sex difference of physiological markers at maximal exercise stress. Women showed significantly lower levels in the cardiopulmonary markers such as maximal systolic blood pressure, ventilation and ventilated lung volume. The maximal heart rate was similar between both sexes, but

Table 6 Changes in swimming speed in backstroke across from 1994 to 2011 for Swiss top 10 swimmers

	Changes (m/s)	Changes (%)	Signifi- cance	Sex differ- ence 1994	Sex differ- ence 2011	Signifi- cance
50-m backstroke						
Women	+0.13	+8.5	P < 0.01	10.58	10.92	P > 0.05
Men	+0.16	+9.4	P < 0.01			
100-m backstroke						
Women	+0.07	+4.8	P < 0.01	9.36	10.01	P = 0.04
Men	+0.12	+7.5	P < 0.01			
200-m backstroke						
Women	+0.06	+4.4	P < 0.01	9.59	10.01	P > 0.05
Men	+0.07	+4.7	P < 0.01			
Changes in swimming speed are presented in m/s and %						

Changes in swimming speed are presented in m/s and %

Licensee OA Publishing London 2013. Creative Commons Attribution License (CC-BY)

FOR CITATION PURPOSES: Kollarz C, Knechtle B, Rüst CA, Rosemann T, Lepers R. Comparison of age of peak swimming speed in elite backstroke swimmers at national and international level. OA Sports Medicine 2013 Oct 01;1(2):19.




Figure 4: Change in age (panels A, C, E) and swimming speed (panels B, D, F) of the international freestyle swimmers for 50 m (panels A and B), 100 m (panels C and D) and 200 m in freestyle (panels E and F) with sex difference across the years.

women performance is limited due to a longer rate-corrected QT interval (time of ventricular de- and repolarisation). The metabolic factors such as blood pH, base excess and blood bicarbonate were similar between men and women with minimal lower levels of blood lactate for men⁴⁰.

These physiological factors seemed to be related to the different levels of muscle power of the sexes. Men have a higher peak leg power compared to women⁴²⁻⁴⁶ mainly due to their higher lean leg volume^{42,44,46}. Seifert

et al.⁴⁷ analysed kinematic changes during a 100-m front crawl sprint for both sexes. High-speed swimmers were characterised by a higher and more stable stroke length. Men swim faster than women because of the greater body height and arm span⁴⁸.

The differences in swimming speed between international and national swimmers can firstly be related to the greater amount of competitors with tighter concentration of the top athletes with top performances. Secondly, international swimmers produce

stability in performances with energetic and biomechanical capacities continuously⁴⁹. Another finding was that the sex difference of national swimmers became greater across the years for 100-m backstroke and all freestyle distances with no changes in international level.

Analysing the change of the sex differences in all swimming events across time, Thiebault et al.¹¹ found a balance in the sex difference in different world records since 1983. With increasing distance, the sex difference became smaller but the levels of sex difference for all swim distances and disciplines are lower than at our data; however, this can be influenced by the greater period of the collected data. Nevill et al.³⁸ analysed the 100-m, 200-m and 400-m freestyle swimming world records from 1957 to 2006 and showed that the sex difference in freestyle remained stable between 8% and 11%. For the 100-m freestyle, the sex difference was at 11%, for the 200-m freestyle it was at 10% and for the 400-m freestyle it was at 9%³⁸. Due to these reasons, a further decline in sex difference might be observed in the future; however, women will not be able to reach the same level as men.

Improvement of swimming speed across years

Our data showed that athletes of both sexes enhanced swimming speed in all distances and disciplines at both national and international levels over the analysed period. The improvement of swim speed throughout years is in agreement with the findings of Berthelot et al.⁵⁰ and Nevill et al.³⁸. Berthelot et al.⁵⁰ analysed data of swimming events from 1963 to 2008, and despite other sports, e.g. track and field, performances in swimming still improved. Nevill et al.³⁸ analysed world records in 100-m, 200-m and 400-m front crawl swimming from 1957 to 2006. They argued that the improvements between the 1950s and 1960s for men

Table 7 Changes in swimming speed in freestyle across from 1994 to 2011 for Swiss top 10 swimmers

	Changes (m/s)	Changes (%)	Significance	Sex difference 1994	Sex difference 2011	Significance
50-m freestyle						
Women	+0.05	+2.8	P < 0.01	10.86	13.30	P < 0.01
Men	+0.11	+4.4	P < 0.01			
100-m freestyle						
Women	+0.04	+2.4	P < 0.01	9.65	12.74	P < 0.01
Men	+0.12	+6.5	P < 0.01			
200-m freestyle						
Women	+0.04	+2.6	P < 0.01	8.77	10.68	P < 0.01
Men	+0.09	+5.3	P < 0.01			
Changes in swimming speed are presented in m/s and %.						

Changes in swimming speed are presented in m/s and %.

and between 1960s and 1970s for women were due to better coaching techniques based on better physiological, biomechanical and psychological knowledge. The impact of the social historical bias in sex difference was great in the early twentieth century. However, biological factors are more important to understand the sex difference today⁵¹.

Berthelot et al.⁵⁰ suggested that age of peak performance might be confounded by introducing a new generation of swimsuits with better hydrodynamic characters. O'Connor et al.⁵² analysed swimming performances of

FINA World Championship in different strokes and distances from 1990 to 2011, with the aim to confirm the influence of technical improvement. Due to the fact that the performances are limited, there were 43 new world records registered at the 2009 World Championship in Rome. At the same time, the full-body polyurethane swimsuit was introduced. O'Connor et al.⁵² claimed the new swimsuit as the main reason for these world records based on the fact that most world records still stand after excluding the full-body swimsuit in 2010. The present data showed significant

improvements in swimming speed across the analysed period. Due to the small period, a plateau was not observed.

Limitations

This cross-sectional data analysis analyses and compares data of national and international swimmers during a period of 17 years. However, it is limited due to lack of determination of body composition^{30,34,35,53}. It is known that they interact with swimming performance⁵⁴. Further studies should include data about body height and weight, body fat and other

Table 8 Changes in swimming speed in freestyle across from 1994 to 2011 for international FINA World Championship swimmers

	Changes (m/s)	Changes (%)	Significance	Sex difference 1994	Sex difference 2011	Significance
50-m freestyle						
Women	+0.06	+3.0	$P = 0.02$	10.64	10.74	P > 0.05
Men	+0.07	+3.1	$P < 0.01$			
100-m freestyle						
Women	+0.05	+2.8	$P < 0.01$	9.67	10.64	P > 0.05
Men	+0.07	+3.5	$P < 0.01$			
200-m freestyle						
Women	+0.03	+1.8	$P = 0.03$	8.77	9.48	P > 0.05
Men	+0.05	+2.7	$P < 0.01$			
Changes in swimming speed are presented in m/s and %.						

Changes in swimming speed are presented in m/s and %.

Licensee OA Publishing London 2013. Creative Commons Attribution License (CC-BY)

FOR CITATION PURPOSES: Kollarz C, Knechtle B, Rüst CA, Rosemann T, Lepers R. Comparison of age of peak swimming speed in elite backstroke swimmers at national and international level. OA Sports Medicine 2013 Oct 01;1(2):19.

anthropometric pattern. Information about nutrition and fluid intake was not recorded, which are known to influence individual performances⁵⁵. Dependent on age and sex, individual demands on nutrition were described^{55,56}. Furthermore, data about training strategies to improve swimming techniques^{47,48,57–61} to optimise recovery phases^{18,62} and warming-up phases^{63,64} which have an effect on swimming performances were not registered. Information about motivation and goal orientation was not recorded. General gender-related differences in motivation⁶⁵, the influence of perfectionism⁶⁶ and mental preparation strategies⁶⁷ are supposed to influence performance.

Conclusion

The present results showed that for freestyle and backstroke and for swimming distance ranging from 50 m to 200 m, elite female swimmers competing at national and international levels achieved peak swimming speed at a younger age (~18–23 years) compared to men (~21–26 years). Across the analysed period, the sex differences in the age of peak swimming speed in backstroke decreased for 50 m and for all distances in freestyle swimming for swimmers competing at national level, while no changes were found for athletes competing at international level. The swimming speed improved over the years in a greater extent for men than for women in all distances and disciplines. Future studies should include data about other swimming styles and the development in a greater variety of distances over a greater analysed period of time.

References

- Bongard V, McDermott AY, Dallal GE, Schaefer EJ. Effects of age and gender on physical performance. *Age (Dordr)*. 2007 Sep;29(2–3):77–85.
- Donato AJ, Tench K, Glueck DH, Seals DR, Eskurza I, Tanaka H. Declines in physiological functional capacity with age: a longitudinal study in peak swimming performance. *J Appl Physiol*. (1985) 2003 Feb;94(2):764–9.
- Tanaka H, Seals DR. Age and gender interactions in physiological functional capacity: insight from swimming performance. *J Appl Physiol*. (1985) 1997 Mar;82(3):846–51.
- Reaburn P, Dascombe B. Endurance performance in masters athletes. *Eur Rev Aging Phys Act*. 2008;5:31–42.
- Lepers R, Knechtle B, Stapley PJ. Trends in triathlon performance: Effects of sex and age. *Sports Med*. 2013 Sep;43(9):851–63.
- Schulz R, Curnow C. Peak performance and age among superathletes: track and field, swimming, baseball, tennis, and golf. *J Gerontol*. 1988 Sep;43(5):P113–20.
- Berthelot G, Len S, Hellard P, Tafflet M, Guillaume M, Vollmer JC, et al. Exponential growth combined with exponential decline explains lifetime performance evolution in individual and human species. *Age (Dordr)*. 2012 Aug;34(4):1001–9.
- Fairbrother JT. Age-related changes in top-ten men's U.S. Masters 50-m freestyle swim times as a function of finishing place. *Percept Mot Skills*. 2007 Dec;105(3 Pt 2):1289–93.
- Leyk D, Erley O, Ridder D, Leurs M, Rütter T, Wunderlich M, et al. Age-related changes in marathon and half-marathon performances. *Int J Sports Med*. 2007 Jun;28(6):513–7.
- Knechtle B, Rüst CA, Rosemann T, Lepers R. Age-related changes in 100-km ultra-marathon running performance. *Age (Dordr)*. 2012 Aug;34(4):1033–45.
- Thibault V, Guillaume M, Berthelot G, El Helou N, Schaal K, Quinquis L, et al. Women and men in sport performance: The gender gap has not evolved since 1983. *J Sports Sci Med*. 2010 June;9(2):214–23.
- Buhl C, Knechtle B, Rüst CA, Rosemann T, Lepers R. A comparison of medley and freestyle performances for national and international swimmers between 1994 and 2011. *Open Access J Sports Med*. 2013 Mar;4:79–87.
- Website: Swiss-swimming: <http://www.rankings.fsn.ch/> [accessed June 2013].
- Website: Swiss-rankings: <http://www.swimrankings.net> [accessed June 2013].
- Website: Federation international de nation: http://www.fina.org/H2O/index.php?option=com_content&view=category&id=271&Itemid=786 [accessed June 2013].
- Website: SAMW (Schweizerische Akademie der medizinischen Wissenschaften): <http://www.samw.ch/en/Ethics/Guidelines/Currently-valid-guidelines.html> [accessed September 2013].
- Guillaume M, Helou El, Nassif H, Berthelot G, Len S, Thibault V, et al. Success in developing regions: World records evolution through a geopolitical prism. *PLoS One*. 2009 Oct;4(10):e7573.
- Issurin VB. New horizons for the methodology and physiology of training periodization. *Sports Med*. 2010 Mar;40(3):189–206.
- Avlonitou E. Somatometric variables for preadolescent swimmers. *J Sports Med Phys Fitness*. 1994 Jun;34(2):185–91.
- Kagawa M, Tahara Y, Moji K, Nakao R, Aoyagi K, Hills AP. Secular changes in growth among Japanese children over 100 years (1900–2000). *Asia Pac J Clin Nutr*. 2011;20(2):180–9.
- Kurokawa N, Nakai K, Suzuki K, Sakurai K, Shimada M, Kameo S, et al. Trends in growth status among schoolchildren in Sendai, Japan, 1994–2003: leveling-off of mean body height and weight. *Tohoku J Exp Med*. 2008 Dec;216(4):371–5.
- Simsek F, Ulukol B, Gulnar SB. The secular trends in height and weight of Turkish school children during 1993–2003. *Child Care Health Dev*. 2005 Jul;31(4):441–7.
- Marmo DB, Zambon MP, Morcillo AM, Guimarey LM. Secular trends of growth of schoolchildren from Paulínia, São Paulo-Brazil (1979/80–1993/94). *Rev Assoc Med Bras*. 2004 Oct–Dec;50(4):386–90.
- Krawczynski M, Walkowiak J, Krzyżaniak A. Secular changes in body height and weight in children and adolescents in Poznań, Poland, between 1880 and 2000. *Acta Paediatr*. 2003;92(3):277–82.
- Ji CY, Hu PJ, He ZH. Secular growth trends in the Chinese urban youth and its implications on public health. *Beijing Da Xue Xue Bao*. 2007 Apr;39(2):126–31.
- Zhen-Wang B, Cheng-Ye J. Secular growth changes in body height and weight in children and adolescents in Shandong, China between 1939 and 2000. *Ann Hum Biol*. 2005 Sep–Oct;32(5):650–65.
- Charles JD, Bejan A. The evolution of speed, size and shape in modern

Licensee OA Publishing London 2013. Creative Commons Attribution License (CC-BY)

FOR CITATION PURPOSES: Kollarz C, Knechtle B, Rüst CA, Rosemann T, Lepers R. Comparison of age of peak swimming speed in elite backstroke swimmers at national and international level. *OA Sports Medicine* 2013 Oct 01;1(2):19.

- athletics. *J Exp Biol.* 2009 Aug;212(Pt 15):2419–25.
28. Geladas ND, Nassis GP, Pavlicevic S. Somatic and physical traits affecting sprint swimming performance in young swimmers. *Int J Sports Med.* 2005 Mar;26(2):139–44.
29. Martin DD, Heckmann C, Jenni OG, Ranke MB, Binder G, Thodberg HH. Metacarpal thickness, width, length and medullary diameter in children--reference curves from the First Zürich Longitudinal Study. *Osteoporos Int.* 2011 May;22(5):1525–36.
30. Wheeler MD. Physical changes of puberty. *Endocrinol Metab Clin North Am.* 1991 Mar;20(1):1–14.
31. Margarey AM, Boulton TJ, Chatterton BE, Schulz C, Nordin BE, Cockington RA. Bone growth from 11 to 17 years: relationship to growth, gender and changes with pubertal status including timing of menarche. *Acta Paediatrica.* 1999 Feb;88(2):139–46.
32. Siders WA, Lukaski HC, Bolonchuk WW. Relationships among swimming performance, body composition and somatotype in competitive collegiate swimmers. *J Sports Med Phys Fitness.* 1993 Jun;33(2):166–71.
33. Tuuri G, Loftin M, Oescher J. Association of swim distance and age with body composition in adult female swimmers. *Med Sci Sports Exerc.* 2002 Dec;34(12):2110–4.
34. Zuniga J, Housh TJ, Mielke M, Hendrix CR, Camic CL, Johnson GO, et al. Gender comparisons of anthropometric characteristics of young sprint swimmers. *J Strength Cond Res.* 2011 Jan;25(1):103–8.
35. Bitar A, Vernet J, Coudert J, Vermorel M. Longitudinal changes in body composition, physical capacities and energy expenditure in boys and girls during the onset of puberty. *Eur J Nutr.* 2000 Aug;39(4):157–63.
36. Lavoie JM, Montpetit RR. Applied physiology of swimming. *Sports Med.* 1986 May–Jun;3(3):165–89.
37. McLean SP, Hinrichs RN. Sex differences in the centre of buoyancy location of competitive swimmers. *J Sports Sci.* 1998 May;16(4):373–83.
38. Nevill AM, Whyte GP, Holder RL, Peyrebrune M. Are there limits to swimming world records? *Int J Sports Med.* 2007 Dec;28(12):1012–7.
39. Vitor Fde M, Böhme MT. Performance of young male swimmers in the 100-meters front crawl. *Pediatr Exerc Sci.* 2010 May;22(2):278–87.
40. Fomin Å, Ahlstrand M, Schill HG, Lund LH, Ståhlberg M, Manouras A, et al. Sex differences in response to maximal exercise stress test in trained adolescents. *BMC Pediatr.* 2012 Aug 20;12:127.
41. Holmér I. Swimming physiology. *Ann Physiol Anthropol.* 1992 May;11(3):269–76.
42. Doré E, Martin R, Ratel S, Duché P, Bedu M, Van Praagh E. Gender differences in peak muscle performance during growth. *Int J Sports Med.* 2005 May;26(4):274–80.
43. Hübner-Woźniak E, Kosmol A, Lutosławska G, Bem EZ. Anaerobic performance of arms and legs in male and female free style wrestlers. *J Sci Med Sport.* 2004 Dec;7(4):473–80.
44. Martin RJ, Dore E, Twisk J, van Praagh E, Hautier CA, Bedu M. Longitudinal changes of maximal short-term peak power in girls and boys during growth. *Med Sci Sports Exerc.* 2004 Mar;36(3):498–503.
45. Nindl BC, Mahar MT, Harman EA, Patton JF. Lower and upper body anaerobic performance in male and female adolescent athletes. *Med Sci Sports Exerc.* 1995 Feb;27(2):235–41.
46. Perez-Gomez J, Rodriguez GV, Ara I, Olmedillas H, Chavarren J, González-Henriquez JJ, Dorado C, Calbet JA. Role of muscle mass on sprint performance: gender differences? *Eur J Appl Physiol.* 2008 Apr;102(6):685–94.
47. Seifert L, Chollet D, Chatard JC. Kinematic changes during a 100-m front crawl: effects of performance level and gender. *Med Sci Sports Exerc.* 2007 Oct;39(10):1784–93.
48. Seifert L, Boulesteix L, Chollet D. Effect of gender on the adaptation of arm coordination in front crawl. *Int J Sports Med.* 2004 Apr;25(3):217–23.
49. Costa MJ, Bragada JA, Mejias JE, Louro H, Marinho DA, Silva AJ, et al. Tracking the performance, energetics and biomechanics of international versus national level swimmers during a competitive season. *Eur J Appl Physiol.* 2012 Mar;112(3):811–20.
50. Berthelot G, Tafflet M, El Helou N, Len S, Escolano S, Guillaume M, et al. Athlete atypicality on the edge of human achievement: performances stagnate after the last peak, in 1988. *PLoS One.* 2010 Jan 20;5(1):e8800.
51. Cheuvront SN, Carter R, Deruisseau KC, Moffatt RJ. Running performance differences between men and women: an update. *Sports Med.* 2005;35(12):1017–24.
52. O'Connor LM, Vozenilek JA. Is it the athlete or the equipment? An analysis of the top swim performances from 1990 to 2010. *J Strength Cond Res.* 2011 Dec;25(12):3239–41.
53. Faulkner JA, Larkin LM, Claflin DR, Brooks SV. Age-related changes in the structure and function of skeletal muscles. *Clin Exp Pharmacol Physiol.* 2007 Nov;34(11):1091–6.
54. Jagomägi G, Jürimäe T. The influence of anthropometrical and flexibility parameters on the results of breaststroke swimming. *Anthropol Anz.* 2005 Jun;63(2):213–9.
55. Stellingwerff T, Maughan RJ, Burke LM. Nutrition for power sports: middle-distance running, track cycling, rowing, canoeing/kayaking, and swimming. *J Sports Sci.* 2011;29(Suppl 1):S79–89.
56. Martínez S, Pasquarelli BN, Romaguera D, Arasa C, Tauler P, Aguiló A. Anthropometric characteristics and nutritional profile of young amateur swimmers. *J Strength Cond Res.* 2011 Apr;25(4):1126–33.
57. Seifert L, Chelensse A, Tourny-Chollet C, Lemaitre F, Chollet D. Effect of breathing pattern on arm coordination symmetry in front crawl. *J Strength Cond Res.* 2008 Sep;22(5):1670–6.
58. Chollet D, Seifert L, Boulesteix L, Carter M. Arm to leg coordination in elite butterfly swimmers. *Int J Sports Med.* 2006 Apr;27(4):322–9.
59. Chollet D, Seifert LM, Carter M. Arm coordination in elite backstroke swimmers. *J Sports Sci.* 2008 May;26(7):675–82.
60. Leblanc H, Seifert L, Baudry L, Chollet D. Arm–leg coordination in flat breaststroke: a comparative study between elite and non-elite swimmers. *Int J Sports Med.* 2005 Nov;26(9):787–97.
61. Takagi H, Sugimoto S, Nishijima N, Wilson B. Differences in stroke phases, arm-leg coordination and velocity fluctuation due to event, gender and performance level in breaststroke. *Sports Biomech.* 2004 Jan;3(1):15–27.

Research study

62. Aspenes ST, Karlsen T. Exercise-training intervention studies in competitive swimming. *Sports Med.* 2012 Jun;42(6):527–43.
63. Balilionis G, Nepocatyč S, Ellis CM, Richardson MT, Neggers YH, Bishop PA. Effects of different types of warm-up on swimming performance, reaction time, and dive distance. *J Strength Cond Res.* 2012 Dec;26(12):3297–303.
64. Arnett MG. Effects of prolonged and reduced warm-ups on diurnal variation in body temperature and swim performance. *J Strength Cond Res.* 2002 May;16(2):256–61.
65. Hanrahan SJ, Cerin E. Gender, level of participation, and type of sport: differences in achievement goal orientation and attributional style. *J Sci Med Sport.* 2009 Jul;12(4):508–12.
66. Stoeber J, Uphill MA, Hotham S. Predicting race performance in triathlon: the role of perfectionism, achievement goals, and personal goal setting. *J Sport Exerc Psychol.* 2009 Apr;31(2):211–45.
67. Houston M, Dolan S, Martin S. The impact of physical, nutritional, and mental preparation on triathlon performance. *J Sports Med Phys Fitness.* 2011 Dec;51(4):583–94.

Licensee OA Publishing London 2013. Creative Commons Attribution License (CC-BY)

FOR CITATION PURPOSES: Kollarz C, Knechtle B, Rüst CA, Rosemann T, Lepers R. Comparison of age of peak swimming speed in elite backstroke swimmers at national and international level. *OA Sports Medicine* 2013 Oct 01;1(2):19.

Competing interests: none declared. Conflict of interests: none declared.
All authors contributed to conception and design, manuscript preparation, read and approved the final manuscript.
All authors abide by the Association for Medical Ethics (AME) ethical rules of disclosure.